

to ED. They wisely note that efficacy should also be assessed by monitoring pharmacodynamic drug effects, such as antimicrobial killing.

1. Keller F. Drug use in the intensive care unit. *Kidney Int* 2012; **82**: 936.
2. Barrantes F, Horwedel T. Drug use and nephrotoxicity in the intensive care unit: are we under-dosing antimicrobials in patients with acute kidney injury with the need of extended dialysis? *Kidney Int* 2012; **82**: 935–936.
3. Perazella MA. Drug use and nephrotoxicity in the intensive care unit. *Kidney Int* 2012; **81**: 1172–1178.
4. Matzke GR, Aronoff GR, Atkinson AJ *et al*. Drug dosing consideration in patients with acute and chronic kidney disease—a clinical update from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int* 2011; **80**: 1122–1137.
5. Mueller BA, Scoville BA. Adding to the armamentarium: antibiotic dosing in extended dialysis. *Clin J Am Soc Nephrol* 2012; **7**: 373–375.

Mark A. Perazella¹

¹Yale University School of Medicine, Section of Nephrology, New Haven, Connecticut, USA

Correspondence: Mark A. Perazella, Yale University School of Medicine, Section of Nephrology, BB 122, 330 Cedar Street, New Haven, Connecticut 06520, USA. E-mail: mark.perazella@yale.edu

Kidney International (2012) **82**, 936–937; doi:10.1038/ki.2012.287

Combining a fruit and vegetable diet with sodium bicarbonate supplementation seems the best dietary option for chronic kidney disease patients

To the Editor: Goraya *et al.*¹ have, in their interesting paper, shown that short-term (30 days) dietary acid reduction due to fruit and vegetable (F + V) or sodium bicarbonate (NaHCO₃) consumption attenuated kidney injury in hypertensive chronic kidney disease (CKD; stadium II) patients undergoing therapy

with angiotensin-converting enzyme inhibitors. The F + V diet reduced systolic blood pressure and body weight, but NaHCO₃ did not affect these parameters. In a comment on this paper, Uribarri and Oh² stated that F + V diets rather than NaHCO₃ supplementation might be the key to halting CKD progression. The truth seems to lie somewhere in between in that simultaneous use of F + V diets and NaHCO₃ is required. In this study, NaHCO₃ caused greater aldosterone reduction than the F + V diet. Because aldosterone is involved in kidney injury, long-term NaHCO₃ supplementation can be more beneficial than the F + V diet alone. Furthermore, the F + V diet reduces urinary sodium concentration. An epidemiological study reported that lower sodium excretion in healthy patients is associated with higher cardiovascular disease mortality.³ Because the Na⁺ cation seems more likely to increase blood pressure when combined with the Cl[−] anion than with HCO₃[−] (ref. 4), providing sodium through bicarbonate rather than kitchen salt seems reasonable. Low long-term dietary compliance is another issue with the F + V diet. Therefore, it seems that CKD patients should use NaHCO₃ supplementation as well as F + V diets.

1. Goraya N, Simoni J, Jo C *et al*. Dietary acid reduction with fruits and vegetables or bicarbonate attenuates kidney injury in patients with a moderately reduced glomerular filtration rate due to hypertensive nephropathy. *Kidney Int* 2012; **81**: 86–93.
2. Uribarri J, Oh MS. The key to halting progression of CKD might be in the produce market, not in the pharmacy. *Kidney Int* 2012; **81**: 7–9.
3. Stolarz-Skrzypek K, Kuznetsova T, Thijs L *et al*. Fatal and nonfatal outcomes, incidence of hypertension, and blood pressure changes in relation to urinary sodium excretion. *JAMA* 2011; **305**: 1777–1785.
4. McCarty MF. Should we restrict chloride rather than sodium? *Med Hypotheses* 2004; **63**: 138–148.

Igor Loniewski¹

¹Sanum Polska, Szczecin, Poland

Correspondence: Igor Loniewski, Sanum Polska, Kurza Stopka 5/C, Szczecin, Zachodniopomorskie 70-535, Poland. E-mail: igorloniewski@sanum.com.pl

Kidney International (2012) **82**, 937; doi:10.1038/ki.2012.289